

Burns and Plastic Surgery Occupational Therapy

Instructions for using Kelocote Spray

Directions for Use

- Spray three times daily after scar massage.
- Shake can well before use.
- Ensure that the affected area is cleaned and then dried.
- Remove the cap from the can and spray along the length of a scar in a sweeping motion.
- Kelocote spray should be applied in a very thin even coat over the whole of the affected area (complete coverage is important for maximum efficacy).
- Kelocote should not pool, drip or run (if it does, you have used too much).
- Allow to dry completely before covering with pressure garments, sun block or make up (Kelocote may stain clothing if not completely dry)

Care of Spray

Kelocote contains flammable gases to help deliver the silicone.

- Extremely flammable, keep away from heat.
- Do not spray on open flame or incandescent material; keep away from sources of ignition.
- Do not smoke whilst using.
- Pressurised container – protect from sunlight and do not expose to temperatures exceeding 50°C/120°F.
- Do not pierce or burn can even after use.
- Do not inhale kelocote spray.
- Do not use near the eyes, nose, mouth or over open wounds.
- For external use only.
- Keep out of reach of children.
- Do not use after the expiry (EXP) date printed on the can.

If you require any further guidance or have any questions about the silicone spray or if the scar area shows any evidence of skin irritation, rash, open or weeping areas, stop using spray and contact your Key Worker

Telephone: (01792) 703980
(Answer Machine available)